

# Persistent high-energy spin excitations in RIXS spectra of optimally doped Bi-2212

*Tuesday, 24 July 2012 20:00 (2 hours)*

M. Guarise, B. Dalla Piazza, H. Ronnow, H. Berger, M. Grioni  
Institute of Condensed Matter Physics, EPFL, CH-1015 Lausanne (Switzerland)

We have exploited high-resolution resonant inelastic x-ray scattering (RIXS) at the Cu L3 edge to measure the full spectrum of spin excitations in Bi-2212. We find that the magnon dispersion and spectral line shape of the insulating Y-substituted parent compound are very similar to those of the AFM insulator Sr<sub>2</sub>CuO<sub>2</sub>Cl<sub>2</sub> [1]. Namely, we observe a dispersion along the magnetic zone boundary, indicative of interactions beyond nearest-neighbors. We also find that clear signatures of spin waves persist in superconducting underdoped and even optimally doped Bi-2212 samples, but develop a remarkable anisotropy. Along the nodal direction the magnon peaks are well-defined and follow the dispersion of the parent insulator. By contrast, only a broad quasielastic tail is seen in the anti-nodal direction. These results suggest a similar anisotropy of the coherence length of the paramagnons.

[1] M. Guarise et al., Phys. Rev. Lett. 105, 157006 (2010)

**Primary author:** GRIONI, Marco (ICMP-EPFL)

**Presenter:** GRIONI, Marco (ICMP-EPFL)

**Session Classification:** Poster Session 2

**Track Classification:** High-Tc Cuprates