

# Dispersive Kondo resonance and hybridized bands in Yb-based compounds

*Monday, 23 July 2012 20:00 (2 hours)*

Masaharu Matsunami, UVSOR Facility  
Tetsuya Hajiri, UVSOR Facility  
Hidetoshi Miyazaki, UVSOR Facility  
Masashi Kosaka, Saitama University  
Shin-ichi Kimura, UVSOR Facility

Heavy-fermion or valence-fluctuation phenomena in strongly-correlated f-electron systems can be derived from the hybridization between conduction band and 4f states (c-f hybridization). Therefore, to directly probe the c-f hybridized electronic structures in the momentum space is important for understanding the heavy-fermion physics. Angle-resolved photoemission spectroscopy (ARPES) is a powerful tool for this purpose. In this work, we have performed ARPES for Yb-based compounds such as YbAl<sub>2</sub>, a prototypical valence fluctuation system, to observe the energy-momentum dispersion curves in the three-dimensional momentum space using a synchrotron radiation.

For YbAl<sub>2</sub>, the c-f hybridized bands and the dispersive Kondo resonance peak have been clearly observed in the APRES spectra. The results are closely associated with the extremely high Kondo temperature (>2000 K). On the basis of these findings, the c-f hybridized electronic structures in Yb compounds will be discussed in comparison with theoretical models such as LDA band calculations and periodic Anderson model.

**Primary author:** MATSUNAMI, Masaharu (UVSOR Facility, Institute for Molecular Science)

**Presenter:** MATSUNAMI, Masaharu (UVSOR Facility, Institute for Molecular Science)

**Session Classification:** Poster Session 1

**Track Classification:** Spin Phenomena