Contribution ID: 122

## Far infrared study of magnetic field-induced normal state of the high temperature superconductor LSCO.

Monday, 23 July 2012 14:00 (25 minutes)

Thomas Timusk, Jungseek Hwang McMaster University, Canada T. Rõõm, U. Nagel, D. Hüvonen, Nat. Inst. Chemical Physics and Biophysics, Estonia D. Hawthorn, S. Wakimoto, H. Zhang, University of Toronto, Canada

We report on the ab-plane optical properties of the magnetic field induced

normal state of La1.94Sr0.06CuO4 (Tc=5.5 K), the first such study. We apply strong

magnetic fields up to 16 T along the c-axis . We find that the magnetic fields, which are strong enough to destroy superconductivity, induce a gap-like depression in the optical conductivity at low frequency along with parallel growth

of a broad absorption peak at higher frequency, just above the gap. The loss of low frequency conductivity in the gap region is in good agreement with dc magneto resistance measurements on samples from the same batch. The

spectral weight loss in the depression at low frequency is

recovered by the spectral weight in the broad peak. Significantly, this spectral weight equals the spectral weight of the superconducting condensate. We interpret our data in terms of a field-induced SDW state that coexists with superconductivity in low fields and grows in amplitude with field in agreement with neutron scattering results of Lake et al. and the phase diagram proposed by Demler et al.

B. Lake et al. Nature, 415, 299, (2002).E. Demler, S, Sachdev and Y. Zhang, PRL, 87, 067202 (2001).

Primary author: TIMUSK, Thomas (McMaster University)

Presenter: TIMUSK, Thomas (McMaster University)

Session Classification: Correlated Physics

Track Classification: Correlated Physics