

Common Fermi-liquid origin of T^2 resistivity and superconductivity in n-type SrTiO₃

Monday, 23 July 2012 10:45 (25 minutes)

Dirk van der Marel, Université de Genève
Dook van Mechelen, Université de Genève
Igor Mazin, Naval Research Laboratories

SrTiO₃ is a semiconductor which, when doped with a low density of electrons, becomes a good conductor with relatively high mobility and strong temperature dependence of the electrical resistivity and the infrared optical conductivity. At low temperatures the material becomes superconducting with T_c below 1 K having a dome-shaped doping dependence, both in the 3D bulk material and at the 2D LaAlO₃/SrTiO₃ interface. The DC resistivity below 100 K has a T^2 temperature dependence. The quasiparticles are in the anti-adiabatic limit with respect to electron-phonon interaction, which renders the interaction mediated through phonons effectively non-retarded. We apply Fermi-liquid theory for the T^2 term in the resistivity, and combine this with expressions for T_c and with the Brinkman-Platzman-Rice (BPR) sum-rule to obtain Landau parameters of n-type SrTiO₃. These parameters are comparable to those of liquid ³He, indicating interesting parallels between these Fermi-liquids despite the differences between the composite fermions from which they are formed. The physics of the doped semiconductor SrTiO₃ stands in stark contrast with the doped cuprates where T_c 's are two orders of magnitude higher and correlate with the T^1 term of the resistivity.

Primary author: VAN DER MAREL, Dirk (Université de Genève)

Presenter: VAN DER MAREL, Dirk (Université de Genève)

Session Classification: Metal-Insulator Systems

Track Classification: Correlated Physics