Contribution ID: 83

α-(BEDT-TTF)2I3: Complex electrodynamic response of the charge-ordered phase

Monday, 23 July 2012 14:45 (12 minutes)

Tomislav Ivek [1], Ivan Kovačević [1], Marko Pinterić [1,2], Bojana Korin-Hamzić [1], Silvia Tomić [1], Conrad Clauss [3], Martin Dressel [3], Dieter Schweitzer [4]

[1] Institut za fiziku, P.O.Box 304, HR-10001 Zagreb, Croatia

[2] Faculty of Civil Engineering, Smetanova 17, 2000 Maribor, Slovenia

[3] 1. Physikalisches Institut, Universität Stuttgart, Pfaffenwaldring 57, D-70550 Stuttgart, Germany

[4] 3. Physikalisches Institut, Universität Stuttgart, Pfaffenwaldring 57, D-70550 Stuttgart, Germany

Much experimental and theoretical attention has been attracted by organic systems with reduced dimensionality and strong Coulomb interactions, and deservedly so due to their novel broken-symmetry phases and corresponding excitations. Here we take a detailed look at the electrodynamics of one of the most prominent charge-ordered systems, the quasi-2D conductor α -(BEDT-TTF)2I3. A semimetal at high temperatures, at 136 K this particular system transitions into an insulating, diamagnetic ground state. Within the insulating phase a long-range commensurate ordering appears in the BEDT-TTF molecular planes, the so-called "horizontal stripe" charge order. [1,2,3] We characterize the charge response of the low-temperature phase using dc resistivity, dielectric and optical spectroscopy in different crystallographic directions within the BEDT-TTF layer. [4,5] Interestingly, two dielectric relaxation modes appear in the kHz-MHz range. The large mode features an anisotropic phason-like behavior, while the small mode presents a soliton-like characteristic. The observed type of excitations agrees with the most relevant physical picture of this charge order as a cooperative bondcharge density wave with ferroelectric-like features. [6] On the other hand, puzzling phenomena including negative differential resistance and voltage oscillations have been reported under application of high electric fields. [7] Our carefully designed electric-field-dependent measurements of conductivity anisotropy within the molecular plane qualitatively confirm these findings; additionally, they reveal novel intriguing behaviors. [8].

[1] Y. Takano et al., J. Phys. Chem. Solids 62, 393 (2001).

[2] Organic Conductors, Special Topics Section, J. Phys. Soc. Jpn. 75, No. 5, (2006).

[3] T. Kakiuchi et al., J. Phys. Soc. Jpn. 76, 113702 (2007).

[4] T. Ivek et al., Phys. Rev. Lett. 104, 206406 (2010)

[5] T. Ivek et al., Phys. Rev. B 83, 165128, (2011).

[6] R. T. Clay et al., J. Phys. Soc. Jpn. 71, 1816 (2002).

[7] K.Tamura et al., J. Appl.Phys.107, 103716(1-5), (2010).

[8] T. Ivek et al., to be submitted (2012).

Primary author: IVEK, Tomislav (Institut za fiziku)

Presenter: IVEK, Tomislav (Institut za fiziku)

Session Classification: Correlated Physics

Track Classification: Correlated Physics