

# Search for Time Reversal Invariance Violation in Resonances of Compound Nuclei Accessible using Epithermal Neutrons

Libertad Barrón Palos\*

Instituto de Física  
Universidad Nacional Autónoma de México

\* for the NOPTREX collaboration



# Matter-antimatter asymmetry in the universe

## Sakharov Criteria


- Conditions out of thermal equilibrium
- Non conservation of baryon number
- C and CP violation ( $CP \longleftrightarrow T$ )

## Searches of TRIV in strongly interacting systems

Complicated theoretical landscape, therefore searches of T violation in different systems, with sensitivity to different possible T-odd mechanisms are necessary:


- Electric Dipole Moments (EDMs)
- T-odd polarized neutron optics

# Electric Dipole Moments



$$\vec{d} = \int \vec{x} \rho(x) d^3x = d \hat{s}$$


# Electric Dipole Moments



$d \neq 0$  constitutes a null test of time reversal invariance  
currently for the neutron  $d_n < 3 \times 10^{-26}$  e·cm


# Hierarchy of CP-odd parameters

[Pospelov & Ritz, Ann. Phys. 318 (2005) 119]



# Hierarchy of CP-odd parameters

[Pospelov & Ritz, Ann. Phys. 318 (2005) 119]



# T violation in p-wave neutron resonances in heavy nuclei

## NOPTREX (Neutron Optics Time Reversal Experiment)

For low-energy neutrons ( $kR \ll 1$ ), the neutron-nucleus interaction can be treated within the framework of neutron optics

$$V_F = \frac{2\pi\hbar^2}{m_n} Nf \quad \leftarrow \text{Fermi potential}$$

with  $m_n$  the neutron mass,  $N$  the number of scattering centers per unit volume and  $f$  the forward scattering amplitude (zero angle elastic scattering)


$$f = a_0 + b_0(\vec{\sigma} \cdot \vec{I}) + c_0(\vec{\sigma} \cdot \vec{k}) + d_0[\vec{\sigma} \cdot (\vec{k} \times \vec{I})]$$

where  $\vec{I}$  is the polarization of the target and  $\vec{\sigma}$  is the neutron spin



The P-odd amplitude [ $c_0(\vec{\sigma} \cdot \vec{k})$ ] has been measured in many nuclear systems, with amplifications by factors of  $10^5$ - $10^6$  due to symmetry mixing in the compound nuclear resonances. [G.E. Mitchell et al. Phys. Rep. 354 (2001) 157]

# $^{139}\text{La}+n$ system

P-odd amplitude



$^{140}\text{La}$  g.s.




[V.W. Yuan et al. Phys. Rev. C 44 (1991) 2187]

# T violation in p-wave neutron resonances in heavy nuclei

- The PV and TRIV effects correspond to the real and imaginary part of the same matrix element (calculated with exactly the same wave functions), therefore it is expected that the TRIV effects would have the same enhancement on resonance as the PV ones.
- Because the neutron optics effects in this case involve elastic scattering at zero angle, the initial and final state coincide and therefore there are no Final State Interactions (FSI) that can mimic T-odd correlations originated from TRIV interactions.
- TRIV effects in neutron transmission on compound nuclei resonances constitute a null test for T invariance.
- The search for TRIV effects in resonances of compound nuclei expands the variety of nuclear systems available ( $^{139}\text{La}$ ,  $^{131}\text{Xe}$ ,  $^{81}\text{Br}$ ,...), helping to provide assurance that possible “accidental” cancelation of TRIV effects due to unknown structural factors in particular systems can be avoided.


# T violation in p-wave neutron resonances in heavy nuclei

In practice...




Measurements of TRIV in this systems is very sensitive to the alignment of the relevant vectors, which are very difficult to control to the required precision.

# Model experiment to search for TRIV in resonances of compound nuclei




[Bowman & Gudkov, Phys. Rev. C 90 (2014) 065503,  
Kabir, Phys. Rev. D 37 (1988) 1856]

# Model experiment to search for TRIV in resonances of compound nuclei



[Bowman & Gudkov, Phys. Rev. C 90 (2014) 065503,  
Kabir, Phys. Rev. D 37 (1988) 1856]

# Model experiment to search for TRIV in resonances of compound nuclei




# T violation in p-wave neutron resonances in heavy nuclei

## TRIV theorem

[Bowman & Gudkov, Phys. Rev. C 90 (2014) 065503]

$$H = \frac{2\pi\hbar^2}{m_n} Nf - \frac{\mu}{2} (\vec{\sigma} \cdot \vec{B})$$

$$f = a_0 + b_0(\vec{\sigma} \cdot \vec{I}) + c_0(\vec{\sigma} \cdot \vec{k}) + d_0[\vec{\sigma} \cdot (\vec{k} \times \vec{I})]$$



If  $d_0=0$  and the apparatus is rotated, with  $\vec{I}$  and  $\vec{B}$  being reversed, then the transmission of neutrons in both states are equal.

Any deviation from the equality in the forward and reversed transmission is a clear manifestation of TRIV interactions (nonzero  $d_0$  coefficient).

# Sensitivity to CP-odd sources: discovery potential

## Model dependent analysis

$$\bar{\lambda} = \frac{\Delta\sigma_{\cancel{P}}}{\Delta\sigma_P} \simeq (-0.47) \left[ \frac{\bar{g}_\pi^{(0)}}{h_\pi^1} + (0.26) \frac{\bar{g}_\pi^{(1)}}{h_\pi^1} \right]$$

$\Delta\sigma_{\cancel{P}}$  P-odd, T-odd asymmetry of interest

$\Delta\sigma_P$  Measured P-odd asymmetry

[Y.H. Song et al., Phys. Rev. C 83 (2011) 065503]

where  $\bar{g}$  and  $h$  are meson-nucleon TRIV and PV coupling constants.

$\bar{g}_\pi^{(0)} < 2.5 \times 10^{-10}$  from the experimental limit on nEDM

$\bar{g}_\pi^{(1)} < 0.5 \times 10^{-11}$  from constraint on the  $^{199}\text{Hg}$  atomic EDM

$h_\pi^1 \sim 4.6 \times 10^{-7}$  from DDH “best value” [DDH, Ann. Phys. 124 (1980) 449]


$$\bar{\lambda} < 10^{-4}$$

$\bar{\lambda}$  can be measured with accuracy of  $10^{-6}$  at existing neutron sources

**Limits on TRIV coupling constants could be improved in neutron optics experiments**

# Sensitivity to CP-odd sources: discovery potential


Estimated for JPARC



Slide by H.M. Shimizu

# Sensitivity to CP-odd sources: discovery potential

Estimated for JPARC



corresponding to  $d_n = 3.0 \times 10^{-26} \text{ e cm}$

corresponding to  $d_n = 3.0 \times 10^{-27} \text{ e cm}$

Slide by H.M. Shimizu

# Complementarity of TRIV searches

- EDMs are ground state properties, while highly excited states in heavy nuclei are involved in neutron optics TRIV effects, which offer a quantitatively different environment with different sensitivity to the many possible sources of CP violation.
- Any single type of TRIV search cannot be equally sensitive to all possible CP violation mechanisms, so it is essential to pursue any experiments in different systems which can be realized with sufficient sensitivity.

# Ongoing and future efforts

## Measurement of $\kappa(J)$ for $^{139}\text{La}$ @JPARC

$$v + iw = \langle \phi_p | V_P + V_{PT} | \phi_s \rangle$$

$$\bar{\lambda} = \frac{\Delta\sigma_{PT}}{\Delta\sigma_P} = \kappa(J) \frac{w}{v}$$


$$\kappa(I + 1/2) = -\frac{3}{2^{2/3}} \left( \frac{2I+1}{2I+3} \right)^{3/2} \left( \frac{3}{\sqrt{2I+3}} \gamma - \sqrt{I} \right)^{-1}$$

$$\kappa(I - 1/2) = -\frac{3}{2^{2/3}} \left( \frac{2I+1}{2I-1} \right) \left( \frac{I}{I+1} \right)^{1/2} \left( -\frac{I-1}{\sqrt{2I-1}} \frac{1}{\gamma} + \sqrt{I+1} \right)^{-1}$$

$$\gamma = \left[ \frac{\Gamma_p^n(I + 1/2)}{\Gamma_p^n(I - 1/2)} \right]^{1/2}$$


Recent  $\gamma$ -ray angular correlation measurements in radiative neutron capture on  $^{139}\text{La}$  yield  $\kappa(J) \sim 1$

[Okudaira et al., Phys. Rev. C 97 (2018) 034622]



# Ongoing and future efforts

Improved precision on  $^{139}\text{La}$  PV asymmetry



- Currently taking place at FP12 of LANSCE
- 1% accuracy goal (currently almost 4%)

# Ongoing and future efforts

## Polarizer for epithermal neutrons

- Epithermal neutron polarizers based on  $^3\text{He}$  spin filters have been developed. Polarizations of the order of 70% can be achieved.

## Targets of heavy nuclei and their polarization

- In the fall of 2018 we plan to perform precision PV asymmetry measurements on  $^{81}\text{Br}$  (0.88 eV resonance) and  $^{131}\text{Xe}$  (3.2 eV resonance), making use of a  $^3\text{He}$  neutron polarizer. Searches for PV effects in other nuclei ( $^{235}\text{U}$  for example) are also planned with this setup.
- A proposal has been submitted to LANSCE to search for P-odd effects in never-measured heavy nuclei (Tl, Ir, Re, W, Ta, Hf, Tm, Er, Ho, Dy, Tb, Gd, Eu, Sm, Om, Nd, Ba, Te, Rh, Mo, Sr, As, Co and Mn) using polarized epithermal neutrons.
- Progress has been made in obtaining polarized  $^{139}\text{La}$  targets (KEK, Kyoto U., PSI). Polarizations of the order of 40% have been achieved in lanthanum aluminate crystals.
- The SIU group is working on considerable improvements in  $^{131}\text{Xe}$  polarized cells.

# Conclusions

- Neutron optics on p-wave resonances in compound nuclei offer a possibility to search for TRIV effects. These effects are free of FSI and constitute a null test for T invariance.
- TRIV effects, just as PV effects, can be enhanced by a factor of  $10^6$  in compound nuclei, due to symmetry mixing on the resonance. This creates a potential for TRIV effects in nuclear resonances to be sensitive enough to improve the constraints on TRIV meson-nucleon couplings.
- TRIV in neutron optics experiment are complementary to EDMs since they constitute quantitatively different systems and can, in principle, access different sources CP violation.
- A complicated theoretical landscape makes it important to perform experiments with different sensitivities to possible CP violation sources.

# NOPTREX collaboration



HIROSHIMA UNIVERSITY



INDIANA UNIVERSITY



# Additional Slides

# Violación de P en resonancias de núcleos compuestos

Table 1  
Parity violations observed by TRIPLE

---

Target

---

$^{81}\text{Br}$   
 $^{93}\text{Nb}$   
 $^{103}\text{Rh}$   
 $^{107}\text{Ag}$   
 $^{109}\text{Ag}$   
 $^{104}\text{Pd}$   
 $^{105}\text{Pd}$   
 $^{106}\text{Pd}$   
 $^{108}\text{Pd}$   
 $^{113}\text{Cd}$   
 $^{115}\text{In}$   
 $^{117}\text{Sn}$   
 $^{121}\text{Sb}$   
 $^{123}\text{Sb}$   
 $^{127}\text{I}$   
 $^{131}\text{Xe}$   
 $^{133}\text{Cs}$   
 $^{139}\text{La}$   
 $^{232}\text{Th}$  below 250 eV  
 $^{232}\text{Th}$  above 250 eV  
 $^{238}\text{U}$


[G.E. Mitchell et al., Physics Reports 354 (2001) 157-241]

# Detector tests



## Detector Prototype

We are designing a current-mode detector for the NOPTREX experiment.

Neutrons hit a  $^{10}\text{B}$  ( $\text{B}_4\text{C}$ ) plate, which has a nice  $1n:1\gamma$  reaction



$\gamma$ -rays are detected using NaI scintillators, shielded in  $^{10}\text{B}$  and lead.



*Slide by D. Schaper*