

Demonstration of Single Barium Ion Sensitivity for Neutrinoless Double Beta Decay Using Single Molecule Fluorescent Imaging

Thursday, May 31, 2018 2:20 PM (20 minutes)

A new method to tag the barium daughter in the double beta decay of ^{136}Xe is described, based on adaptation of the single molecule fluorescent imaging (SMFI) technique. Individual barium dications chelated on a transparent plate are detected at a significance of 12.9σ , with a spatial resolution of 2 nm rms. Observation of a single-step photo-bleach transition confirms the interpretation of single ion sensitivity. This result is the first step toward an essentially background-free technique in the search for neutrino-less double beta decay in ^{136}Xe , based on robust event discrimination by SMFI in a high-pressure xenon gas TPC.

E-mail

nygren@uta.edu

Collaboration name

NEXT Collaboration

Primary authors: Mr MCDONALD, Austin (University of Texas at Arlington); Prof. JONES, Ben (University of Texas at Arlington); Prof. NYGREN, David (University of Texas at Arlington)

Presenters: Prof. NYGREN, David (University of Texas at Arlington); Dr NYGREN, David (Lawrence Berkeley National Laboratory)

Session Classification: Neutrino Masses and Neutrino Mixing

Track Classification: NMNM