

# Demonstration of Single Barium Ion Sensitivity for Neutrinoless Double Beta Decay Using Single Molecule Fluorescent Imaging

*Thursday, 31 May 2018 14:20 (20 minutes)*

A new method to tag the barium daughter in the double beta decay of  $^{136}\text{Xe}$  is described, based on adaptation of the single molecule fluorescent imaging (SMFI) technique. Individual barium dications chelated on a transparent plate are detected at a significance of  $12.9\sigma$ , with a spatial resolution of 2 nm rms. Observation of a single-step photo-bleach transition confirms the interpretation of single ion sensitivity. This result is the first step toward an essentially background-free technique in the search for neutrino-less double beta decay in  $^{136}\text{Xe}$ , based on robust event discrimination by SMFI in a high-pressure xenon gas TPC.

## E-mail

nygren@uta.edu

## Collaboration name

NEXT Collaboration

**Primary authors:** Mr MCDONALD, Austin (University of Texas at Arlington); Prof. JONES, Ben (University of Texas at Arlington); Prof. NYGREN, David (University of Texas at Arlington)

**Presenters:** Dr NYGREN, David (Lawrence Berkeley National Laboratory); Prof. NYGREN, David (University of Texas at Arlington)

**Session Classification:** Neutrino Masses and Neutrino Mixing

**Track Classification:** NMNM