

## Measurement of Polarization Observables in the Reaction $\gamma p \rightarrow K^+ \Lambda$

*Tuesday, 29 May 2018 18:10 (20 minutes)*

Spin observables are important to understand the production mechanisms of hyperons, as well as the contribution of intermediate baryon resonances.  $\Lambda$  polarization observables have been studied extensively in the recent decades using the reaction  $\gamma + p \rightarrow K^+ + \Lambda$ . This talk presents the measurement of transferred polarization coefficients  $C_x$  and  $C_z$ , and the induced polarization  $P$ , using a new set of high statistics data, obtained using the CEBAF Large Acceptance Spectrometer (CLAS) detector at Jefferson Lab. The photon beam energy range is 1.117 to 5.45 GeV. These results ( $C_x$ ,  $C_z$  and  $P$ ) are extracted simultaneously using the Maximum Likelihood Method. The measurements for  $C_x$  and  $C_z$  have nearly an order of magnitude increase in events compared to previously published results and also extend the kinematic range for  $W > 2.54$  GeV, important for both the search for high-mass nucleon states as well as to provide information about non-resonant contributions.

### **E-mail**

sadhi003@fiu.edu

**Primary author:** ADHIKARI, Shankar (Florida International University)

**Presenter:** ADHIKARI, Shankar (Florida International University)

**Session Classification:** QCD, Hadron Spectroscopy, and Exotics

**Track Classification:** QCDHS