

The Nucleon Axial Coupling from Quantum Chromodynamics

Thursday, 31 May 2018 16:10 (30 minutes)

The *axial coupling of the nucleon*, g_A , is the strength of its coupling to the *weak axial current* of the Standard Model, much as the electric charge is the strength of the coupling to the electromagnetic current. This axial coupling dictates, for example, the rate of β -decay of neutrons to protons and the strength of the attractive long-range force between nucleons. Precision tests of the Standard Model in nuclear environments require a quantitative understanding of nuclear physics rooted in Quantum Chromodynamics, a pillar of this theory. The prominence of g_A makes it a benchmark quantity to determine theoretically, a difficult task as the theory is non-perturbative. Lattice QCD provides a rigorous, non-perturbative definition of the theory which can be numerically implemented. In order to determine g_A , the lattice QCD community has identified two challenges which must be overcome to achieve a 2% precision by 2020: the excited state contamination must be controlled and the statistical precision must be markedly improved. Here we report a calculation of $g_A^{\text{QCD}} = 1.271 \pm 0.013$ using an unconventional method that overcomes these challenges.

E-mail

chiachang@lbl.gov

Collaboration name

CalLat

Funding source

DOE

Primary author: Dr CHANG, Chia Cheng (LBL)

Presenter: Dr CHANG, Chia Cheng (LBL)

Session Classification: Nuclear Forces and Structure, NN Correlations, and Medium Effects

Track Classification: NFS