

# Nucleon Form Factors in Dispersively Improved Chiral Effective Theory

*Saturday, June 2, 2018 6:10 PM (20 minutes)*

We present a new method for calculating the nucleon electromagnetic form factors (EM FFs) combining Chiral Effective Field Theory (ChEFT) and dispersion analysis [1]. The FFs are expressed as dispersive integrals over the two-pion cut at timelike  $t > 4M_\pi^2$ . The spectral functions are computed using elastic unitarity, chiral pion-nucleon amplitudes (LO, NLO, partial N2LO), and timelike pion FF data. The method effectively includes  $\pi\pi$  rescattering effects and the  $\rho$  resonance and leads to major improvements compared to traditional ChEFT calculations. Higher-mass isovector and isoscalar  $t$ -channel states are described by effective poles, whose strength is fixed by sum rules (charges, radii). We obtain excellent agreement with the spacelike proton and neutron FF data up to  $Q^2 \sim 1 \text{ GeV}^2$ . We predict the values of the higher FF derivatives with minimal uncertainties and study their collective behavior (multiple dynamical scales, unnatural sizes). Our approach provides a FF parametrization with proper analyticity and theoretical uncertainty estimates, which can be used for analysis of low- $Q^2$  elastic scattering data and extraction of the proton radius [2].

[1] J.M. Alarcon, C. Weiss, Phys. Rev. C 97, 055203 (2018).

[2] J.M. Alarcon, C. Weiss, arXiv:1803.09748 [hep-ph].

## E-mail

weiss@jlab.org

**Primary author:** WEISS, Christian (Jefferson Lab)

**Presenter:** WEISS, Christian (Jefferson Lab)

**Session Classification:** QCDHS / PPHI

**Track Classification:** QCDHS