

## Recent Developments in the Proton Radius Puzzle

*Tuesday, 29 May 2018 09:15 (35 minutes)*

The CODATA-recommended value of the proton rms charge radius  $r_p$  is based on hydrogen/deuterium spectroscopy and electron-proton/deuteron scattering experiments. Since 2010, a discrepancy between the  $r_p$  value determined from muonic hydrogen spectroscopy and the CODATA-recommended value has led to intense research activity both in theory and experiment. The situation has recently become even more puzzling as a contradiction appeared between two new hydrogen spectroscopy results: the 2S-4P transition frequency [Beyer *et al.*, *Science* 358, 2017] agrees with the muonic  $r_p$  value, while the 1S-3S transition frequency presented here supports the CODATA value.

In our experiment at Laboratoire Kastler Brossel (Paris, France), the two-photon 1S-3S hydrogen transition is excited in an atomic beam, with a continuous-wave 205-nm laser obtained by sum frequency generation in a non-linear crystal. The transition frequency is measured with respect to the LNE-SYRTE Cs clock by means of a frequency comb. The second-order Doppler shift and other systematic effects have been included in the analysis to determine the 1S-3S transition frequency with a relative uncertainty of  $9 \times 10^{-13}$  [Fleurbaey *et al.*, PRL accepted for publication, 2018]. The value of the proton radius which can be inferred is in very good agreement with the CODATA-recommended value.

In this talk, I will present the current situation of the proton radius puzzle and our latest results.

### E-mail

helene.fleurbaey@nist.gov

**Primary author:** FLEURBAEY, Helene (NIST)

**Presenter:** FLEURBAEY, Helene (NIST)

**Session Classification:** Plenary 1

**Track Classification:** QCDHS