



Emerging Detection Capabilities for Nuclear Deterrence

Keegan J. Kelly



2025 Workshop for Applied Nuclear Data Activities

Outline


- White-Source ($n,2n$) and ($n,3n$) Measurements Capabilities with CLYC-7 n - n Coincidences
- The DAPPER array
- GENESIS Forward Analysis at the LBNL 88-inch Spectrometer
- Exotic fission measurements with the FRIB High-Rigidity Spectrometer
- Microcalorimeters for Nuclear Data Measurements



Fusion Reactors Rely on $(n,2n)$ and $(n,3n)$ Rxns





ITER_D_3FM52L - Radiation environment for equipment during operations by R Juarez



- n breeding essential for t production via $^6\text{Li}(n,t)$, to drive $d-t$ – $^9\text{Be}(n,2n)$ and $^{208}\text{Pb}(n,2n)$
- Activation-based flux measurements motivate a suite of $(n,2n)$ measurements

Traditionally: γ -rays, Activation, or n Counting

Calculated with CoH₃ - T. Kawano, Springer Proceedings in Physics 254 (2021) 27



→ Traditional methods do not measure emitted n information

→ Detection of both ($n, 2n$) neutrons captures 100% of strength.

Continuous white-source neutron measurements are ideal, but neutron TOF degeneracies are problematic



Degeneracies of White Sources can be Solved

Neutron energies for $(n,2n)$ for $(n,3n)$ reactions at white sources are degenerate


$$T_n^{(1)} = t_n^{inc} + t_n^{(1)} = \text{First TOF}$$

$$T_n^{(2)} = t_n^{inc} + t_n^{(2)} = \text{Second TOF}$$

BUT, measuring one neutron energy breaks the degeneracy!

$$t_n^{(1)} = \left(\frac{l_{out}}{c} \right) \left[1 - \left(1 + \frac{E_n^{(1)}}{m_n c^2} \right)^{-2} \right]^{-1/2} \Rightarrow t_n^{inc} = T_n^{(1)} - t_n^{(1)} \Rightarrow E_n^{inc}, E_n^{(2)}$$

LANSCCE can provide continuous $(n,2n)$ and $(n,3n)$ measurements with emitted neutron energy and angular information



Degeneracies of White Sources can be Solved

Neutron energies for $(n,2n)$ for $(n,3n)$ reactions at white sources are degenerate


$$T_n^{(1)} = t_n^{inc} + t_n^{(1)} = \text{First TOF}$$

$$T_n^{(2)} = t_n^{inc} + t_n^{(2)} = \text{Second TOF}$$

BUT, measuring one neutron energy breaks the degeneracy!


$$t_n^{(1)} = \left(\frac{l_{out}}{c} \right) \left[1 - \left(1 + \frac{E_n^{(1)}}{m_n c^2} \right)^{-2} \right]^{-1/2} \Rightarrow t_n^{inc} = T_n^{(1)} - t_n^{(1)} \Rightarrow E_n^{inc}, E_n^{(2)}$$

LANSCCE can provide continuous $(n,2n)$ and $(n,3n)$ measurements with emitted neutron energy and angular information



Accomplished with CLYC-7 E_n Data and CoGNAC

- Upgrade CoGNAC to include a series of high-volume CLYC-7 scintillators
- $^{35}\text{Cl}(n,p)$ measures $E_n^{(1)}$ directly
- EJ-309 and CLYC-6 detectors provide $T_n^{(2)}$ measurement to low energy




Applicable to 20+ $(n,2n)$ and $(n,3n)$ measurements for DOE SC NP FES, and could lead to a decade+ campaign for OES / SAT and PAT

Funded by the DOE Early Career Research Program





GENESIS is performing simultaneous energy differential and integral measurements of the (n,xny) reactions to help better inform evaluation


26 neutron detectors $N(E_n, \theta_n)$




7 HPGe Detector $N(E_n, \theta_n)$



Neutron flux measurement with $E_{max} < 22$ MeV



Reaction model parameters are iteratively adjusted to best reproduce GENESIS observables



Detector Array for Photons, Protons, and Exotic Residues

Contact: Alan McIntosh; alanmcintosh[at]tamu.edu


DAPPER

Neutron capture x-sect needed on unstable isotopes for



- stockpile science
- advanced reactor design
- nuclear forensics
- nuclear astrophysics

Can't measure everything. Can't measure certain things at all.

Need for nuclear data to constrain models, improve predictions.



- photons: 800 lbs BaF2 → individual gamma energy, total gamma energy, gamma multiplicity; segmented for Doppler correction
 - protons: S3 annular silicon → excitation energy
 - exotic residues: zero degree ionization chamber → discriminate beam & reaction of interest @ >600k pps
-
- Multiple analysis methods: Forward (multi-step cascade), Oslo, Shape
 - Systematic measurements along isotopic chains
 - Multiple detector arrays: DAPPER, Hyperion, etc.
 - Multiple surrogate reactions: (d,p), (p,p'), (3He,4He), (t,p), (p,d)...



NNSA: DE-NA0003841
DE-NA0004150 (CENTAUR)
DOE-NP: DE-FG02-93ER40773



TEXAS A&M UNIVERSITY

Cyclotron Institute




U.S. DEPARTMENT OF
ENERGY



LA-UR-25-21178


Detector Array for Photons, Protons, and Exotic Residues

Contact: Alan McIntosh; alanmcintosh [at] tamu.edu




High efficiency: 24% @ 8 MeV

clustering (adback)
boosts efficiency



Excitation vs gamma energy:
Large yield in total detection

- E* and E_g contain information on
- Nuclear Level Density
 - Photon Strength Function



Extracted 58Fe PSF and NLD
Parallel analyses agree


55Fe PSF and NLD in progress
See Arthur Alvarez's Poster!



NNSA: DE-NA0003841
DE-NA0004150 (CENTAUR)
DOE-NP: DE-FG02-93ER40773



Overview of HRS Consisting of HRS-HTBL & HRS-SPS




R. Zegers - HRS for Fission Measurements - FIESTA 2024

- Pure FRIB beam is incident on a production target
 - Select isotope of interest, and transport to HRS
- Interaction with Reaction Target produces nucleus that will fission “in flight”



Multiple Operational Modes of HRS-SPS Facilitates Various Science Programs under Optimal Conditions

- Multiple operational modes available in a single hardware setup
 - Lattice layout is optimized for **high-resolution mode** in which the majority of experiments will run
 - Neutron-invariant-mass mode** is specialized for invariant-mass spectroscopy with fast neutron detection at forward angles, with reaction target placed in front of the first dipole
- Interface with auxiliary detectors is coordinated with user community
 - Input from users (organized through the HRS Working Group) is continuously sought to ensure that the HRS design accommodates user program and facilitates detector installation










R. Zegers - HRS for Fission Measurements - FIESTA 2024

- Excitation energy of fissioning nucleus can be measured event-by-event
- High rigidities and excitations from surrogate reactions emulate neutron capture preceding fission





Adaptability to Measure Variety of Rxn Outputs



- Able to fission fragment and prompt observable data directly with CGMF, FREYA, etc.
 - Models limited by structure of neutron-rich fragments from fission



Microcalorimeters as Specialized Tools for ND




- For narrow-focus, specialized cases, microcalorimeters are unrivaled
- Commonly used for decay or x-ray spectroscopy



Images from J. Ward, LA-UR-23-32260



LA-UR-25-21178



μ Cals May Yield Elusive Actinide Scattering Data



The onset of the $^{239}\text{Pu}(n,n')$ reaction to the 7.9 keV first excited state is a fundamental missing component for nuclear model development


- The trajectory of $\sigma(n,n')$ can vary wildly based on details of first inelastic channel

μ Cals can observe both (n,n) and (n,n') using only the recoiling nucleus



μ Cals May Yield Angular Distributions Too!

Fig. Courtesy of J. Surbrook



The energy deposited into the μ Cal is correlated to the energy and angle of the emitted neutron

→ Neutron angular distributions can be calculated from measurements of the parent nucleus




14 MeV Measurements with DT Generators


Fig. Courtesy of

J. Surbrook


Incident neutron



Elastic scatter



Inelastic scatter



*High-precision 14 MeV Measurements for fusion
and more applications are possible*

See poster by Jason Surbrook, and Lee Bernstein talk with μ Cals



The Berkeley team is also developing a DT-API system on campus to measure $(n_{14}, x n \gamma)$ cross sections with little-to-no flux uncertainty

Simple Count Rate Estimate

$$R_{\alpha-\gamma/n} = \sigma(N)_{sample} \Phi_n$$


For a 10 g sample 1 m away from a 1 cm scintillator @ 5 cm:

$$N_{sample} \approx 10^{23} \text{ atoms}$$

$$\Phi_n \approx 10^8 \frac{n}{s} \times \frac{1}{100} \rightarrow \approx 10^9 \frac{n}{day}$$

$$\rightarrow R_{\alpha-\gamma/n} = \frac{10^8}{\sigma \text{ (barn)} \cdot \text{day}}$$

10⁵ peak counts/day for a 0.1% detection efficiency



See L. Bernstein talk tomorrow in Fusion Nuclear Data session

THANK YOU!

- CoGNAC (n, xn) detector development funded by DOE Office of Science via Early Career Research Program.
- μ Cal initial investigations for nuclear data funded by NA-113, as part of the CoGNAC neutron scattering program at LANL.
- GENESIS and DT-API work at LBNL were performed under the auspices of the Office of Nonproliferation Research and Development (NNSA/NA-22) and the US Nuclear Data Program at Lawrence Berkeley National Laboratory (Contract No. DEAC02-05CH11231) and the Stewardship Science Academic Alliance Program under Grant DE-NA0004064 at the University of California, Berkeley.

Direct questions to kkelly@lanl.gov

