



Contribution ID: 151

Type: Oral

Spectroscopy of Trans-fermium Nuclei Using the Argonne Gas-Filled Analyzer

Monday, 13 June 2022 11:30 (20 minutes)

Spectroscopy of Trans-fermium Nuclei Using the Argonne Gas-Filled Analyzer

D. Seweryniak,¹ T. Huang,^{1,2} K. Auranen,^{1,11} A.D. Ayangeakaa,^{1,12} B.B. Back,¹ P. Bender,³ M.P. Carpenter,¹ P. Chowdhury,³ R.M. Clark,⁴ P. Copp,¹ Z. Favier,¹⁰ K. Hauschild,⁸ X.-T. He,⁵ R.D. Herzberg,⁶ D. Ho,³ H. Jayatissa,¹ T.L. Khoo,¹ F.G. Kondev,¹ G. Morgan,⁷ C. Morse,⁴ A. Korichi,⁸ T. Lauritsen,¹ J. Li,¹ C. Mueller-Gatermann,¹ D. Potterveld,¹ W. Reviol,¹ A. Rogers,³ S. Saha,³ G. Savard,¹ K. Sharma,³ S. Stolze,¹ S. Waniganeththi,³ G. Wilson,¹ J. Wu,¹ Y.-F. Xu,⁵ S. Zhu⁹

¹Argonne National Laboratory, Argonne, IL 60439, USA

²Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

³University of Massachusetts, Lowell, MA 01854, USA

⁴Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

⁵College of Materials Science and Technology, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China

⁶University of Liverpool, Liverpool, UK

⁷Louisiana State University, Baton Rouge, LA 70803, USA

⁸CSNSM, IN2P3-CNRS Orsay, France

⁹Brookhaven National Laboratory, Upton, NY 11973, USA

¹⁰CEA, Département de Physique Nucléaire, Université Paris-Saclay, 91191, Gif-sur-Yvette, France

¹¹Accelerator Laboratory, Department of Physics, University of Jyväskylä, FI-40014 Jyväskylä, Finland

¹²Triangle Universities Nuclear Laboratory, Duke University, Durham, North Carolina 27708, USA

Deformed nuclei near the $Z=100$, $N=152$ deformed shell gaps are a stringent testing ground for nuclear models which are used to describe the heaviest known nuclei. Nuclei in this region have been studied using in-beam, K -isomer, α -decay and spontaneous fission spectroscopic methods. To extend these studies to odd- A , odd-odd, and to heavier nuclei the Argonne Gas-Filled Analyzer (AGFA) was constructed. During the talk, recent decay and isomer spectroscopy experiments with AGFA in stand-alone mode and in-beam spectroscopy experiments with AGFA coupled to the Gammasphere γ -ray detector array will be reviewed. Among others, the observation of the ground-state rotational band in the fissile nucleus ^{254}Rf and the discovery of the new isotope ^{251}Lr will be discussed. The moment of inertia deduced for ^{254}Rf shades light on the shape evolution in this region while the observed α -decay fine structure in ^{251}Lr provides information on single-proton orbitals near the Fermi surface. Plans for experimental program with AGFA will be also presented.

This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under contract number DE-AC02-06CH11357. This research used resources of ANL's ATLAS facility, which is a DOE Office of Science User Facility.

Primary author: SEWERYNIAK, D. (Physics Division, Argonne National Laboratory, Argonne, IL 60439)

Presenter: SEWERYNIAK, D. (Physics Division, Argonne National Laboratory, Argonne, IL 60439)

Session Classification: NS2022 Plenary

Track Classification: Oral Presentations