



Contribution ID: 131

Type: Oral

## Investigation of the $^{46}\text{Ar}$ proton wave function: the $^{46}\text{Ar}(^3\text{He}, d)^{47}\text{K}$ direct reaction

*Tuesday 14 June 2022 11:50 (20 minutes)*

Direct reactions represent a unique mechanism to investigate the nuclear-structure properties and the nature of single-particle states of nuclei along shell closures. In this contribution, the  $^{46}\text{Ar}(^3\text{He}, d)^{47}\text{K}$  proton-transfer reaction is proposed for the study of properties of the ground state of the radioactive neutron-rich  $^{46}\text{Ar}$  isotope. The interest behind this isotope stems from the observed discrepancies between the well-established shell model with SDPF-U interaction and measurements of transition probability  $B(E2)$  between the ground state and the first excited state. The proton component of the wave function has been pointed out as the source of this discrepancy [1,2].

The measurement, performed at the SPIRAL 1 facility in GANIL, France with a post-accelerated radioactive 9 MeV/u  $^{46}\text{Ar}$  beam impinging on a high-density cryogenic  $^3\text{He}$  target [3], aimed at quantifying the transfer cross section to the  $3/2^+$  level relative to the  $1/2^+$  ground state in  $^{47}\text{K}$ . The experiment relied on a state-of-the-art experimental setup for a precise reconstruction of the kinematics of the reaction. The heavy reaction fragment was identified by the high acceptance magnetic spectrometer, VAMOS [4], while the high-granularity silicon DSSD detector, MUGAST [5], allowed the measurement of the angular distribution of the light ejectile while also performing particle identification. The AGATA gamma-ray tracking germanium array [6] measured the photons produced by the decay of the  $^{47}\text{K}$  excited states.

The experimental evidence indicates a substantially suppressed  $L=2$  transfer to the first excited state of  $^{47}\text{K}$ , at odds with shell-model calculations that predict the  $s_{1/2}$  and  $d_{3/2}$  orbitals as almost degenerate and not entirely occupied. The results will be discussed in the framework of *ab initio* and mean-field calculations. In these theoretical results, the low occupancy of the  $s_{1/2}$  orbital, in agreement with the high relative spectroscopic factor measured, implies a central depletion of the proton wavefunction. This work presents the first experimental evidence of this phenomenon in  $^{46}\text{Ar}$ .

- [1] A. Gade et al., Phys. Rev. C 74, 034322 (2006)
- [2] S. Calinescu et al., Phys. Rev. C 93, 044333 (2016)
- [3] F. Galtarossa et al., Nucl. Inst. Meth. A 1018, 165830 (2021)
- [4] M. Rejmund et al., Nucl. Inst. Meth. A 646, 184-191 (2012)
- [5] M. Assié et al., Nucl. Inst. Meth. A 1014, 165743 (2021)
- [6] S. Akkoyun et al., Nucl. Inst. Meth. A 668, 26-58 (2012)

**Authors:** BRUGNARA, Daniele (Laboratori Nazionali di Legnaro, INFN, Italy); GOTTARDO, Andrea (INFN LNL); Dr ASSIÉ, Marlène (IJCLab, France); Dr MENGONI, Daniele (INFN Padova, University of Padova)

**Presenter:** BRUGNARA, Daniele (Laboratori Nazionali di Legnaro, INFN, Italy)

**Session Classification:** NS2022 Plenary

**Track Classification:** Oral Presentations