



Contribution ID: 114

Type: Oral

## Beta decay strength of $^{78}\text{Ni}$ to neutron-unbound states revealed by $^{79}\text{Cu}$

Thursday, 16 June 2022 16:10 (20 minutes)

$^{78}\text{Ni}$  is one of the few radioactive doubly-magic nuclei on the nuclear landscape [1]. It has a large neutron-to-proton ratio and engages in beta-delayed neutron and gamma emissions. Decays of Cu isotopes with significant neutron excess provide a laboratory to study such doubly-magic nuclei. The nuclear shell structure of  $^{79}\text{Cu}$  ( $N=50$ ;  $Z=29$ ) is highly similar to the one of  $^{78}\text{Ni}$  ( $N=50$ ;  $Z=28$ ), except an extra proton above the  $Z=28$  shell gap in  $1f_{5/2}$  orbital [2]. The similarities in the nuclear shell structure are also reflected in the beta decay properties.  $^{79}\text{Cu}$  is also a strong beta-delayed neutron emitter, which allows us to infer beta-strength function ( $S_{\beta}(E)$ ) to neutron-unbound states in the beta decay of  $^{78}\text{Ni}$ .

The determination of ( $S_{\beta}(E)$ ) requires measurement of the intensity of neutron resonances post beta decay. Neutron-gamma coincidences are also vital to arranging strength with excitation energy. An experiment for measuring the strength distribution in the  $^{78}\text{Ni}$  region ( $27 \geq Z \geq 33$ ) was performed at the Radioactive Ion Beam Factory (RIBF) facility at RIKEN Nishina Center, JAPAN using a YSO-based implantation detector [3] and VANDLE [4] array for Time-of-Flight-based spectroscopy of the beta-delayed neutrons. For  $\gamma$ -ray detection, two HPGe clovers and 10  $LaBr_3$  detectors were used.

This work reports the first direct measurements of beta-decay strength to neutron-unbound states in the decay of  $^{81,80,79}\text{Cu}$  crossing over the  $N=50$  shell closure. The results from the experiment are compared to NuShellX [5] calculations with various sets of single-particle energies and residual interactions. Finally,  $^{78}\text{Ni}$  beta decay is predicted based on the model best describing  $^{79}\text{Cu}$ . The results reveal neutron and proton spin-orbit partners providing significant contributions to ( $S_{\beta}(E)$ ) in the beta decay of  $^{78}\text{Ni}$ . The results also reveal the impact of the  $N=50$  on the shell gap in the decay of  $^{80,81}\text{Cu}$ .

### References

1. R. Taniuchi *et al.*, Nature 569, 53–58 (2019).
2. L. Olivier *et al.*, Phys. Rev. Lett. 119, 192501 (2017).
2. M. Singh *et al.*, (drafted).
4. W.A. Peters *et al.*, Nucl. Instrum. Methods Phys. A 836, 122 (2016).
5. NuShellX, W.D.M. Rae, <http://www.garsington.eclipse.co.uk>.

**Primary authors:** SINGH, Maninder (University of Tennessee, Knoxville); Prof. GRZYWACZ, Robert; Dr YOKOYAMA, Rin; COX, Ian; Dr ZHENGYU, Xu

**Presenter:** SINGH, Maninder (University of Tennessee, Knoxville)

**Session Classification:** NS2022 Plenary

**Track Classification:** Oral Presentations