



Contribution ID: 82

Type: Oral

Study of intruder states towards ^{78}Ni with lifetime measurements following $^{82}\text{Se}(d,p)^{83}\text{Se}$

Quadrupole interaction between protons and neutrons drives the nucleus into deformed configurations at low excitation energies. Around the $N=50$ shell gap, intruder states with spins $1/2^+$ and $5/2^+$, originating from the $s_{1/2}$ and $d_{5/2}$ orbitals were first observed in ^{83}Se and, later on, in the other $N=49$ isotones ^{87}Kr , ^{81}Ge and ^{79}Zn . In ^{83}Se these intruder states reach energies of around 500 keV, the lowest among the other $N=49$ isotones. The ^{83}Se nucleus is at the mid of the proton fp -shell and it should have the maximum of quadrupole correlations which makes it a good candidate to understand the collectivity of the particle-hole intruder states in this region, lowered in energy by large quadrupole correlations. Indeed, large-scale shell model calculations predict a quenching of the energy of the intruder states in ^{83}Se , at variance with the experimental data [1]. Lifetime measurements of the intruder states of ^{83}Se would give an indication about their wave function and would allow estimating the degree of the $N=50$ core breaking in the ground state of Se isotopes. Moreover, such measurements could shed light on the behavior of the $N=50$ shell gap towards ^{78}Ni , a double-magic nucleus in which intruder configurations competing in energy with the spherical ones have also been found [2].

We will report on the results obtained from a recent experiment performed in Laboratori Nazionali di Legnaro, where lifetimes of the intruder-state band were measured using Recoil Distance Doppler-Shift Method and Doppler-Shift Attenuation Method. A beam of ^{82}Se , with intensity 0.02 pnA, accelerated at 270 MeV by the ALPI-TANDEM accelerator at LNL-INFN, impinged into a deuterated polyethylene (C_2D_4) target which was evaporated on a 6 mg/cm^2 thick gold layer. The GALILEO γ -array was coupled to the SPIDER silicon-array, allowing to obtain the needed channel selectivity through coincidence measurements between γ rays and protons coming from the (d,p) transfer reaction. The results on lifetimes will be discussed in the framework of large-scale shell-model calculations and mean-field approaches, pointing out the role of the collectivity of low-lying intruder configurations.

[1] C. Wraith et al., Nature, 569, 53-58 (2019).

[2] R. Taniuchi et al., Phys. Lett. B. 771, 385-391 (2017).

Primary authors: PELLUMAJ, Julgen (INFN-LNL); GOTTARDO, Andrea (INFN LNL)

Presenter: PELLUMAJ, Julgen (INFN-LNL)

Session Classification: Poster Session

Track Classification: Poster Presentations