



Contribution ID: 152

Type: Oral

Constraining the $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ reaction rate using the GADGET II TPC

The $^{15}\text{O}(\alpha,\gamma)^{19}\text{Ne}$ breakout reaction from the hot CNO cycles is significant to the thermonuclear runaway that causes type I X-ray bursts on accreting neutron stars. At breakout temperatures ($\approx 0.5\text{GK}$), this reaction is strongly dominated by a single resonance with center of mass energy 506 Kev corresponding to a ^{19}Ne state having excitation energy of 4.03 MeV. An experimental upper limit has been placed on its strength, but the lower limit on the resonance strength and therefore the astrophysical reaction rate is unconstrained experimentally. Since the lifetime of the state is well known, only a finite experimental value for the alpha-particle branching ratio is needed to determine the rate. With this strong motivation, we have proposed an experiment to measure a finite value for the branching ratio and hence the reaction rate at FRIB using an upgraded version of the Gaseous Detector with Germanium Tagging (GADGET) and this proposal has been accepted by the FRIB's first Program Advisory Committee (PAC). These measurements will proceed via the $^{20}\text{Mg}(\beta\alpha)^{15}\text{O}$ decay sequence using the β decay of ^{20}Mg followed by a proton emission to the 4.03 MeV ^{19}Ne state. These decay events yield a characteristic signature: the emission of a proton and alpha particle. To achieve the high granularity necessary for the identification of this characteristic signature, we have upgraded the GADGET's Proton Detector into a time projection chamber (TPC). A MICROMEGAS board with 1024 ($2.2 \times 2.2\text{ mm}^2$) pads and high-density GET electronics has been installed to accommodate the large number of electronics channels. In order to test the functionality as a TPC, a ^{228}Th source is placed inside the gas handling system of the detector which allows us to bleed ^{220}Rn α inside the TPC. The α tracks from the ^{220}Rn decay have been successfully seen, demonstrating the system's functionality. This upgraded version is known as GADGET II and has three distinct elements: the Segmented Germanium Array (SeGA) for γ - detection, a beam-pipe cross that houses a beam energy degrader and diagnostics, and the new TPC. The TPC has been simulated using the ATTPCROOTv2 data analysis framework based on the FairRoot package for ^{20}Mg and ^{220}Rn decay events. Based on these simulations, a machine learning algorithm is being developed that will be integrated with the ATTPCROOTv2 analysis framework to identify candidates in the data. Transfer learning will be used to refine the machine-learned models after the experiment using real in - situ data on $^{20}\text{Mg}(\beta\alpha)^{19}\text{Ne}$ events for single protons and $^{20}\text{Na}(\beta\alpha)^{16}\text{O}$ daughter-decay events for single alphas. These simulations will be useful in selecting events of interest based on their unique signature to provide background free measurements.

This work has been supported by the U. S. Department of Energy under award no: DE-SC0016052 and the U. S. National Science Foundation under award no: 1565546 and 1913554.

Primary authors: MAHAJAN, Ruchi (Facility for Rare Isotope Beams, Michigan State University); ADAMS, A. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); ALLMOND, J. (Oak Ridge National Laboratory, Oak Ridge, Tennessee 37830, USA); ALVAREZ-POL, H. (IGFAE, Universidade de Santiago de Compostela, E-15782 Santiago de Compostela, Spain); ARGO, E. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); AYYAD, Y. (IGFAE, Universidade de Santiago de Compostela, E-15782 Santiago de Compostela, Spain); BARDAYAN, D. (Department of Physics, University of Notre Dame, Notre Dame, Indiana 46556, USA); BAZIN, D. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); BUDNER, T. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); CHEN, A. (Department of Physics and Astronomy, McMaster University,

Canada); CHIPPS, K. (Oak Ridge National Laboratory, Oak Ridge, Tennessee 37830, USA); DAVIDS, B. (TRIUMF, Vancouver, British Columbia, Canada); DOPFER, J. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); FRIEDMAN, M. (The RACAH institute of Physics, Hebrew University of Jerusalem, Israel); FYNBO, H. (Aarhus University, Aarhus C, Denmark); GRZYWACZ, R. (University of Tennessee, Knoxville, USA); JOSE, J. (Universitat Politècnica de Catalunya, Barcelona, Spain); LIANG, J. (Department of Physics and Astronomy, McMaster University, Canada); PAIN, S. (Oak Ridge National Laboratory, Oak Ridge, Tennessee 37830, USA); PEREZ-LOUREIRO, D. (Canadian Nuclear Laboratories, Canada); POLLACCO, E. (IRFU, CEA Saclay, Gif-sur-Ivette, France); PSALTIS, A. (Technical University of Darmstadt, Germany); RAVISHANKAR, S. (Department of Computational Mathematics, Science and Engineering, Michigan State University, USA); ROGERS, A. (University of Massachusetts Lowell, USA); SCHAEDIG, L. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); SUN, L.J. (School of Physics and Astronomy, Shanghai Jiao Tong University, China); SURBROOK, J. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); WHEELER, T. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); WEGHORN, L. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA); WREDE, C. (Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA)

Presenter: MAHAJAN, Ruchi (Facility for Rare Isotope Beams, Michigan State University)

Session Classification: Poster Session

Track Classification: Poster Presentations