

# Recent constraints on axion-photon and axion-electron couplings with the CAST experiment

*Thursday, September 12, 2013 5:20 PM (20 minutes)*

“Axions are DM candidates that can be produced in the solar core. The CERN Axion Solar Telescope (CAST) is a helioscope looking for axions coming from the solar core to Earth. The experiment, located at Geneva (Switzerland) is able to follow the Sun during sunrise and sunset. Four x-ray detectors are mounted on both ends of the magnet waiting for a photon from axion-to-photon conversion due to the Primakoff effect. With the completion of Phases I and II, CAST has been able to scan for axions with masses up to 1.16eV. We will present recent constraints on the axion-photon coupling.

Recently, non-minimal axion models are also receiving increasing attention. For instance, WD cooling hypothesis relies solely on the axion-electron coupling. Non-hadronic axion models, have a tree-level axion-electron interaction: the Sun produces a strong axion flux by bremsstrahlung, Compton scattering, and axio-recombination, the “BCA processes.”Based on a new calculation of this flux, including for the first time axio-recombination, we will also present limits on the axion-electron Yukawa coupling  $g_{ae}$  and axion-photon interaction strength  $g_{a\gamma}$  using the CAST phase-I data (vacuum phase).”

**Primary author:** RUZ, Jaime (CERN)

**Presenter:** RUZ, Jaime (CERN)

**Session Classification:** Dark Matter VIII

**Track Classification:** Dark Matter